

February 1, 2008

LBNL-53265

July 2003

Production of axial-vector D_{sJ} in e^+e^- annihilation

Mahiko Suzuki

Department of Physics and Lawrence Berkeley National Laboratory

University of California, Berkeley, California 94720

(February 1, 2008)

Abstract

If one of the recently discovered charmed-strange mesons ($D_{sJ}(2317)$) is the 0^+ state of $c\bar{s}$, the other ($D_{sJ}(2460)$) is most likely the 1^+ state with $j = \frac{1}{2}$. They could be produced in e^+e^- annihilation at $E_{e^+e^-}^{\text{cm}} = m_{\Upsilon(4S)}$ either by fragmentation from $c\bar{c}$ jets or as decay products of the B mesons from $\Upsilon(4S) \rightarrow B\bar{B}$. If one analyzes the $c\bar{c}$ jet events and the $\Upsilon(4S)$ decay events separately, one will have a direct test as to whether $D_{sJ}(2460)$ is the $j = \frac{1}{2}$ state or not, how much $D_{sJ}(2460)$ is mixed with the $j = \frac{3}{2}$ state, and also whether the four-quark interpretation is viable.

PACS number(s) 13.25.Ft, 13.20.He, 12.38.Bx, 12.39.Hg

arXiv:hep-ph/0307118v1 8 Jul 2003

Typeset using REVTeX

I. INTRODUCTION

The BaBar Collaboration [1] discovered a narrow peak at 2317 MeV in the invariant mass of $D_s\pi^0$. The decay into $D_s\pi^0$ suggests $J^P = 0^+$ for this state $D_{sJ}(2317)$ since good candidates already exist for 1^- and 2^+ . The observation of this peak was subsequently confirmed by the CLEO [2] and the Belle Collaboration.

The BaBar data show another sharp peak at 2460 MeV in the invariant mass of $D_s^*\pi^0$ ($D_s\gamma\pi^0$). Although BaBar was initially cautious in calling it as a resonance, CLEO concluded that it is another narrow resonance. Absence of the decay $D_{sJ}(2460) \not\rightarrow D_s\pi^0$ leads us to speculate that $D_{sJ}(2460)$ is one of $J^P = 1^+$ states and forms with $D_{sJ}(2317)$ a $j = \frac{1}{2}$ multiplet in terms of total momentum $\mathbf{j} = \mathbf{l} + \mathbf{s}$ of the $\bar{s}$ quark in the heavy-light limit of $c\bar{s}$. Then the existing 1^+ resonance $D_{s1}(2536)$ [3] should be assigned to a $j = 3/2$ multiplet with the 2^+ candidate $D_{sJ}(2573)$.

Theorists have worked on spectroscopy of the heavy-light quark states for long time. Schnitzer [4] argued on the basis of the l - s coupling in a naive two-body potential model that for heavy-light mesons, the $j = \frac{3}{2}$ multiplet should be lighter than the $j = \frac{1}{2}$ multiplet. It was pointed out many years later that K meson resonances seemed to show this “inversion” of the spin-orbit coupling sign despite the relative lightness of the s -quark [5].¹ Then the spin-orbit inversion in the heavy-light system was studied systematically by Isgur [6]. More recently a detailed computation was presented in the potential model [7]. However, the potential picture has an uncertainty of long-distance physics since the energy scale of the heavy-light potential is the reduced mass, *i.e.*, the light mass, not by the heavy mass. Upon the discovery of $D_{sJ}(2317)$, Cahn and Jackson [8] re-examined the $c\bar{s}$ states with the potential model. It is fair to say that the potential model is inconclusive about which of the j -multiplets is heavier than the other. By keeping the inversion scenario back in mind, many theorists proposed the exotic possibility that $D_{sJ}(2317)$ is a four-quark state or its mixing [9,10]. A cursory examination by lattice QCD was also reported [11].

Another approach to the heavy-light mesons is to treat the light component as a chiral-symmetric cloud instead of a constituent quark [12]. Bardeen *et al* made a strong case for noninversion of $j = \frac{1}{2}$ and $j = \frac{3}{2}$ with detailed calculations [13]. Among others the observed D_{sJ} mass difference, $m(2460) - m(2317) \simeq m(1^-) - m(0^-) = 143$ MeV, is a successful consequence of chiral symmetry. As for magnitude of the splitting between $j = \frac{1}{2}$ and $\frac{3}{2}$, however, an earlier work [14] had found much larger uncertainty than Bardeen *et al.* did.

While the case for $j = \frac{1}{2}$ appears strong for $D_{sJ}(2317)$ and $D_{sJ}(2460)$, an independent experimental confirmation is desirable. At $E_{e^+e^-}^{(\text{cm})} = \text{the } \Upsilon(4S) \text{ mass}$, D_{sJ} can be produced through $c\bar{c}$ jets or B decays. Two types of processes occur with roughly the same rate and can be separated without difficulty by event topology and by B decay vertices. We point out here that production rates of the axial-vector D_{sJ} in $c\bar{c} \rightarrow D_{sJ}X$ and in $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow D_{sJ}X$ are sensitive to $j = \frac{1}{2}$ vs $\frac{3}{2}$ and therefore that analyzing the $D_{sJ}(2460)$ production in $c\bar{c}$ and in $B\bar{B}$ separately will give us an additional clue about as to whether $D_{sJ}(2460)$ is $j = \frac{1}{2}$ or

¹Since then, however, the measured branching fractions of $\tau \rightarrow K_1\nu_\tau$ had shifted so that the inversion argument is no longer supported by τ decay data. Only the s - d ratio of $K_1 \rightarrow K^*\pi/\rho K$ may favor the inversion, if at all.

$\frac{3}{2}$ of $c\bar{s}$, or else a four-quark state $c\bar{s}q\bar{q}$. The two types of production occur with roughly the same rate and separable by event topology and B decay vertices.

II. B DECAY

Production of charmed strange mesons is one of the dominant nonleptonic B decay processes. It occurs mainly through the effective decay operators of tree type:

$$\begin{aligned}\mathcal{O}_1 &= (\bar{c}s)_{V-A}(\bar{b}c)_{V-A}, \\ \mathcal{O}_2 &= (\bar{c}c)_{V-A}(\bar{b}s)_{V-A},\end{aligned}\tag{1}$$

where colors are contracted within each bracket. Although detailed comparison between theory and experiment has not been available for $B \rightarrow D_s$, the similar decays $B \rightarrow \bar{D}\pi$, $\bar{D}^*\pi$, $\bar{D}\rho$ and so forth that occur through $(\bar{u}d)_{V-A}(\bar{b}c)_{V-A}$ and $(\bar{u}c)_{V-A}(\bar{b}d)_{V-A}$ have been measured with good accuracy [3]. These color-favored two-body decays agree well with the theoretical values computed in the factorization approximation. The factorization is even simpler for the inclusive color-favored decays since they are free from the quark distribution involving the spectator quark. We therefore proceed by assuming that the color-favored decay $B \rightarrow D_{sJ}X$ is described by the factorization.² In the factorization limit only D_{sJ} of 0^- , 1^- , and 1^+ can be produced. As for the 1^+ states, we shall see below that the 1^+ state of $j = \frac{1}{2}$ would be produced preferentially for $m_s/m_c \ll 1$.

In the simple factorization the inclusive decay $B \rightarrow D_{sJ}X$ is determined by the short-distance quark decay of $\bar{b} \rightarrow \bar{c}(c\bar{s})$. While inclusive production of 0^- and 1^- in raw data contains the contribution from strong and electromagnetic cascade decays of higher D_{sJ} states, production of 1^+ is very likely free of such contamination. The reason is that only the radially excited 0^- and $1^\pm$ states below the DK threshold are possible sources of cascade decays down to the 1^+ states in the factorization. Such excited states have not been seen in experiment. They are expected to be above the DK threshold. Therefore the $D_{s1}(2460)$ reconstructed in B decay may be counted entirely as primary decay products of B meson.

The production amplitude for $\bar{b} \rightarrow 1^+\bar{c}$ is given by

$$A(\bar{b} \rightarrow D_{s1j}\bar{c}) = (G_\mu/\sqrt{2})V_{cb}^*V_{cs}(C_1 + C_2/3)f_{Aj}m_{D_{s1j}}\epsilon^\mu(\bar{v}_b\gamma_\mu\gamma_5v_c), \quad (j = 1/2, 3/2) \tag{2}$$

where the axial-vector decay constant f_{Aj} is defined with the normalization $\langle \mathbf{p}|\mathbf{p}' \rangle = (2\pi)^3 2E_{\mathbf{p}}\delta(\mathbf{p} - \mathbf{p}')$ by

$$\langle D_{s1j}(p, \epsilon) | (\bar{c}\gamma^\mu\gamma_5s) | 0 \rangle = f_{Aj}m_{D_{s1j}}\epsilon^\mu. \tag{3}$$

We have introduced the additional subscript j for D_{s1} to distinguish between two eigenstates of $\mathbf{j}^2$. If one evaluates Eq. (3) in the rest frame of D_{s1j} by treating D_{s1j} as being made of the c -quark and the remainder carrying the $\bar{s}$ -quark quantum numbers (still denoted by $\bar{s}$), the left-hand side is written in the Pauli spinors as

²This is not true for the color-suppressed decays such as $B^0 \rightarrow \bar{D}^0\pi^0$ and charmonium production. The factorization-forbidden charmonia of $J^{PC} = 0^{++}$ and 2^{++} are abundantly produced. [15]

$$\begin{aligned} \langle D_{s1j}(\mathbf{0}, \epsilon, j) | (\bar{c} \gamma \gamma_5 s) | 0 \rangle = N \left[\left(\frac{1}{E_s + m_s} + \frac{1}{E_c + m_c} \right) i \chi_c^\dagger (\mathbf{p}_s \times \boldsymbol{\sigma}) \chi_{\bar{s}} \right. \\ \left. + \left(\frac{1}{E_s + m_s} - \frac{1}{E_c + m_c} \right) \chi_c^\dagger \mathbf{p}_s \chi_{\bar{s}} \right], \end{aligned} \quad (4)$$

where N is an normalization factor. The spinors form 3P_1 and 1P_1 in the first and the second term, respectively, in the right-hand side of Eq. (4). In the limit of $m_s/m_c \rightarrow 0$, the right-hand side approaches $\chi^\dagger \boldsymbol{\sigma} (\boldsymbol{\sigma} \cdot \mathbf{p}_s) \chi_{\bar{s}}$ up to an overall constant, which is exactly the $j = \frac{1}{2}$ combination of 3P_1 and 1P_1 . Therefore, the weak axial-vector current can produce only the $j = \frac{1}{2}$ state of D_{s1} in the large m_c limit [16]. For $m_s/m_c \neq 0$, the weak current produces the combination of

$$|D_{s1\frac{1}{2}}\rangle \cos \alpha - |D_{s1\frac{3}{2}}\rangle \sin \alpha, \quad (5)$$

where

$$\tan \alpha = \frac{2\sqrt{2}}{3(E_c + m_c)/(E_s + m_s) + 1} \quad (6)$$

in the phase convention of $|j = \frac{1}{2}\rangle = \sqrt{\frac{1}{3}}|^1P_1\rangle - \sqrt{\frac{2}{3}}|^3P_1\rangle$ and $|j = \frac{3}{2}\rangle = \sqrt{\frac{2}{3}}|^1P_1\rangle + \sqrt{\frac{1}{3}}|^3P_1\rangle$. For a nonrelativistic binding with $m_s/m_c \simeq \frac{1}{3}$, the mixture of $|j = \frac{3}{2}\rangle$ is small ($\tan^2 \alpha \simeq 0.09$ in probability) and production of $D_{s1\frac{3}{2}}$ is almost negligible. To obtain the production rates of the mass eigenstates, one needs to know about a small mixing between $j = \frac{1}{2}$ and $\frac{3}{2}$ in the mass eigenstates:

$$\begin{aligned} |D_{s1}(2460)\rangle &= \cos \theta |D_{s1\frac{1}{2}}\rangle - \sin \theta |D_{s1\frac{3}{2}}\rangle, \\ |D_{s1}(2536)\rangle &= \sin \theta |D_{s1\frac{1}{2}}\rangle + \cos \theta |D_{s1\frac{3}{2}}\rangle. \end{aligned} \quad (7)$$

Then the ratio of the branching fractions for two mass eigenstates is

$$\frac{\mathcal{B}(B \rightarrow D_{s1}(2536)X)}{\mathcal{B}(B \rightarrow D_{s1}(2460)X)} = \tan^2(\alpha - \theta). \quad (8)$$

In the case that $D_{s1}(2460)$ consists mostly of $j = \frac{1}{2}$, the mixing angle θ is $O(m_s/m_c)$ so that the production rate of $D_{s1}(2536)$ is one order of magnitude smaller than that of $D_{s1}(2460)$. Theoretical estimate of the value of θ is not possible because of unknown long-distance effects. We should use Eq. (8) to determine the mixing angle θ albeit Eq. (6) has some model dependence.

If the value of the decay constant f_{Aj} is given for $j = \frac{1}{2}$, we can compute the branching fraction of the inclusive $B \rightarrow D_{s1\frac{1}{2}}X$ decay. The authors in [16] gave one estimate, which corresponds to $f_A(\equiv f_{A\frac{1}{2}}) \simeq 200\text{MeV}$ for $m_c \simeq 1.35\text{GeV}$ in our definition Eq. (3). Using this value in the amplitude of Eq. (2), we obtain with a straightforward computation the branching fraction,

$$\begin{aligned} \mathcal{B}(B \rightarrow D_{s1\frac{1}{2}}X) &= \frac{\mathcal{B}(B \rightarrow X l^+ \nu_l)_{exp}}{\Gamma(B \rightarrow X l^+ \nu_l)_{th}} \times \Gamma(B \rightarrow D_{s1\frac{1}{2}}X), \\ &\simeq (1.7 \times 10^{-2}) \times (f_A/200\text{MeV})^2, \end{aligned} \quad (9)$$

where $\mathcal{B}(B \rightarrow Xl^+\nu_l)_{\text{exp}} \simeq 0.104$ [3], $C_1 + \frac{1}{3}C_2 \simeq 1.02$ [17], and the short-distance corrected value of $\Gamma(B \rightarrow Xl^+\nu_l)_{\text{th}}$ have been used to obtain the numerical result. This number is subject to uncertainty of the values for $m_b(\simeq 4.5\text{GeV})$ and $m_c(\simeq 1.35\text{GeV})$. The branching fraction is quite large since it is one of the dominant factorizable processes. If we compare two-body decays $B \rightarrow D_{s1\frac{1}{2}}\overline{D}$ and $D_s\overline{D}$, for example, we find

$$\mathcal{B}(B \rightarrow D_{s1\frac{1}{2}}\overline{D})/\mathcal{B}(B \rightarrow D_s\overline{D}) \simeq 1.5 \times (f_A/f_{D_s})^2, \quad (10)$$

where f_{D_s} is the decay constants of D_s .

III. FRAGMENTATION FROM CHARM-ANTICHARM JET

Fragmentation of a heavy meson from a heavy quark was studied by many theorists in perturbative pictures [18]. In the heavy limit of a heavy quark, the fragmentation functions for 1P_1 and 3P_1 have a similar dependence on $z = 2E/m_b$ and ratio of the integrated fragmentation probabilities is $\simeq 2/3$ in the perturbative calculation [19] when the $^3P_1 - ^1P_1$ mass splitting is ignored. In terms of the ratio of $j = \frac{1}{2}$ to $\frac{3}{2}$, this number corresponds to $\simeq 0.88$. After the $O(m_s/m_c)$ corrections are included, the ratio shifts a little but stays close to unity: $\int D_{j=\frac{1}{2}}(z)dz / \int D_{j=\frac{3}{2}}(z)dz \simeq 1$. Actually, physics of fragmentation of a heavy-light meson is not entirely a short-distance process even if one takes the heavy quark limit. We do not know of how to estimate nonperturbative effects reliably. One can understand complexity of long-distance effects if one thinks of cascade feeding from higher resonance states.

For two 1^+ states, the orbital wavefunctions are the same in nonrelativistic models. Cascade contributions are unimportant since higher resonances decay into DK channels. If dominant long-distance effects are spin independent like the confining force, the *ratio* of the fragmentation probabilities would not change much with long-distance effects. In the absence of a compelling reason for otherwise, it is not unreasonable to expect that nonperturbative effects do not upset the perturbative prediction on the fragmentation ratio:

$$\mathcal{B}(e^+e^- \rightarrow c\overline{c} \rightarrow D_{s1\frac{3}{2}}X) \simeq \mathcal{B}(e^+e^- \rightarrow c\overline{c} \rightarrow D_{s1\frac{1}{2}}X). \quad (11)$$

This is markedly different from B decay in which $D_{s1\frac{1}{2}}$ is dominantly produced.

IV. FOUR-QUARK STATE

Many theorists proposed [9] that $D_{sJ}(2317)$ may be an 0^+ state of $c\overline{s}q\overline{q}$. Although it has been speculated that some of light scalar mesons might be four-quark states [20], we have not yet had a resonance that is proven to be a four-quark meson. Consequently, we do not have much knowledge of dynamical properties of four-quark states such as production and decay.

We consider four-quark states, which we denote them generically by $D_s^{(4)}$. First in B decay. The production amplitude for $D_s^{(4)}$ is obtained by superposition of a four-quark production amplitude in momentum space:

$$A_{B \rightarrow D_s^{(4)} X}(p_X) = \prod_{i=1,2,3} \int \frac{d^3 \mathbf{q}_i}{(2\pi)^3} \tilde{\Psi}(\mathbf{q}_i) A_{\bar{b} \rightarrow (c\bar{s}q\bar{q})\bar{c}}(p_X, \mathbf{q}_i), \quad (12)$$

where $\tilde{\Psi}(\mathbf{q}_i)$ is the four-quark wavefunction in momentum space, $\mathbf{q}_i$ ($i = 1, 2, 3$) are the relative quark momenta inside $D_s^{(4)}$, and p_X is momentum of $D_s^{(4)}$. In the loose-binding approximation, the decay amplitude turns into a simple form,

$$A_{B \rightarrow D_s^{(4)} X}(p_X) \simeq \Psi(\mathbf{0}) A_{b \rightarrow (c\bar{s}q\bar{q})\bar{c}}(p_X, \mathbf{0}), \quad (13)$$

where $\Psi(\mathbf{0})$ is the four-quark wavefunction in coordinate space with all three relative coordinates set equal to zero. $|\Psi(\mathbf{r}_j)|^2$ has dimension of the ninth power of energy. For a loosely bound molecular DK state,

$$|\Psi(\mathbf{0})|^2 = O(\Lambda_{QCD}^6 \Delta^3) \quad (14)$$

where Δ is the binding energy. For intrinsic four-quark states in which no $q\bar{q}$ pair is in a color-singlet, $|\Psi(\mathbf{0})|^2$ would be comparable with or smaller than that of the molecular state since the binding is loose. As for the production amplitude, the relevant effective interactions are six-quark operators. The dominant interaction is of the form

$$\mathcal{L}_{int} \sim (G_\mu/\sqrt{2}) V_{cb}^* V_{cs} (\pi\alpha_s/E^3) (\bar{b}\gamma_\mu(1-\gamma_5)c) (\bar{c}q\bar{q}s)^\mu, \quad (15)$$

where $(\bar{c}q\bar{q}s)^\mu$ is a Lorentz vector made of four quark fields such as $(\bar{c}\gamma^\mu\partial_\nu q)(\bar{q}\gamma^\nu s)$, and E is determined by the energy scale involved in creation of q and $\bar{q}$. Then simple dimension counting gives us

$$\Gamma(B \rightarrow D_s^{(4)} X) \sim G_\mu^2 |V_{cb}^* V_{cs}|^2 (\pi\alpha_s)^2 \left| \frac{\Psi(\mathbf{0})}{\sqrt{m_{D_s^{(4)}}}} \right|^2 \frac{m_B^3}{E^6}. \quad (16)$$

Using Eq. (14), we can estimate the branching fraction. If we express in the ratio to the branching to $B \rightarrow D_s X$ for comparison,

$$\mathcal{B}(B \rightarrow D_s^{(4)} X) \simeq (\pi\alpha_s)^2 \left(\frac{\Lambda_{QCD}}{E} \right)^6 \left(\frac{\Delta^3}{m_{D_s^{(4)}} f_{D_s}^2} \right) \times \mathcal{B}(B \rightarrow D_s X). \quad (17)$$

Since the four-quark binding energy Δ is much smaller than $m_{D_s^{(4)}}$ and f_{D_s} , $\mathcal{B}(B \rightarrow D_s^{(4)} X)$ is minuscule as compared with $\mathcal{B}(B \rightarrow D_s X)$. The same estimate as Eq. (17) holds for perturbative fragmentation of $D_s^{(4)}$ from $c\bar{c}$ jets. We thus conclude that short-distance production of $D_s^{(4)}$ is negligibly small both in B decay and in jet fragmentation. In fact, suppression of a loosely bound multiparticle state is a general rule at high energies. The factor $\Delta^3/m_{D_s^{(4)}} f_{D_s}^2$ in Eq. (17) comes from the ratio of the wavefunctions of $D_s^{(4)}$ and D_s . However, there is a chance of large long-distance enhancement in the final state if $D_s^{(4)}$ and a $c\bar{s}$ meson state of the same quantum numbers happen to be almost degenerate in mass. In this case the transition of $D_{sJ}(c\bar{s}) \leftrightarrow D_s^{(4)}$ is enhanced by the factor $|\Psi(\mathbf{0})|^2/(M_{D_s^{(4)}} - M_{D_{sJ}})^2$. This enhancement factor is square of the D_{sJ} - $D_s^{(4)}$ mixing itself. We may state therefore that production of $D_s^{(4)}$ is possible only if it has a large mixing to a quark-antiquark meson state [10].

How much could $D_s(2317)$ be produced if it is the 0^+ state of $D_s^{(4)}$ mixed with a $c\bar{s}$ state? Since $D_s(c\bar{s})$ of $J^P = 0^+$ cannot be produced in the factorization, a mixing to it does not help production of $D_s^{(4)}$ in B decay. On the other hand $D_s^{(4)}$ production through fragmentation would be realized if it is mixed substantially with $D_{sJ}(c\bar{s})$ of $J^P = 0^+$. Since the ratio of fragmentation functions of 0^+ and 1^+ is approximately 0.36 for $m_s/m_c \simeq 1/3$ in the perturbative calculation [19], even a 50-50 mixing to $D_{s0}(c\bar{s})$ would allow fragmentation of $D_s^{(4)}$ only at the level of one fifth of $D_{s1\frac{1}{2}}$. If $D_s^{(4)}$ of $J^P = 1^+$ mixes with $D_{s1\frac{1}{2}}$ strongly, we may be able to see it in B decay. In fragmentation, mixing to 1^+ of either $j = \frac{1}{2}$ or $j = \frac{3}{2}$ helps $D_s^{(4)}$ production.

V. SUMMARY

The BaBar, CLEO, and Belle Collaborations should be able to sort out $\Upsilon(4S)$ events and $c\bar{c}$ jet events by event topology and by B meson decay vertices. By analyzing the two types of events separately, we shall obtain useful information as to which of $D_{s1}(2460)$ and $D_{s1}(2536)$ is $j = \frac{1}{2}$ or $j = \frac{3}{2}$: The 1^+ meson produced abundantly in B decay is $D_{s1\frac{1}{2}}$. The $D_{s1\frac{3}{2}}$ meson should be looked for in the $c\bar{c}$ jet events. If both D_{s1j} states should happen to be produced significantly in B decay, or if only one of two D_{s1j} states is produced from the $c\bar{c}$ jet, it would be an indication of a large mixing between $j = \frac{1}{2}$ and $\frac{3}{2}$, that is, failure of the heavy quark approximation to the c quark.

ACKNOWLEDGMENTS

This work is supported in part by the Director, Office of Science, Office of High Energy and Nuclear Physics, Division of High Energy Physics, of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-0098840.

REFERENCES

- [1] BaBar Collaboration, B. Aubert *et al.*, hep-ex/0304021.
- [2] CLEO Collaboration, D. Besson *et al.*, hep-ex/0305100.
- [3] *Review of Particle Physics*, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [4] H.J. Schnitzer, Phys. Lett. **76B**, 461 (1978); Nucl. Phys. **B207**, 131 (1982).
- [5] M. Suzuki, Phys. Rev. D **50**, 4708 (1994); D **51** 1428(E), (1995).
- [6] N. Isgur, Phys. Rev. D **57**, 4041 (1997).
- [7] M. DiPierro and E. Eichten, Phys. Rev. D **64**, 114004 (2001).
- [8] R.N. Cahn and J.D. Jackson, hep-ph/0305012.
- [9] T. Barnes, F.E. Close, and H.J. Lipkin, hep-ph/0305025; H-Y. Chen and W-S. Hou, hep-ph/0305038; E. van Beveren and G. Rupp, hep-ph/0305035; K. Terasaki, hep-ph/0305213.
- [10] T.E. Browder, S. Pakvasa, and A.A. Petrov, hep-ph/0307054.
- [11] G.S. Bali, hep-ph/0305209.
- [12] H. Cheng, C. Cheung, G. Lin, T. Yan, and H. Yu, Phys. Rev. D **46**, 5060 (1992); G. Burdman and J.F. Donoghue, Phys. Lett. **B280**, 287 (1992); W.A. Bardeen and C.T. Hill, Phys. Rev. D **49**, 409 (1994); A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli, and A.D. Polosa, Phys. Rev. D **58**, 034004 (1998); A. Hiorta and J.O. Eeg, Phys. Rev. D **66**, 074001 (2002).
- [13] W. A. Bardeen, E.J. Eichten, and C.T. Hill, hep-ph/0305049.
- [14] A. Deandrea, N. Di Bartolomeo, R. Gatto, G. Nardulli, and A.D. Polosa, Phys. Rev. D **58**, 034004 (1998).
- [15] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **88**, 031802 (2002); **89**, 011803 (2002).
- [16] P. Colangelo, G. Nardulli, and N. Paver, Phys. Lett. **B293**, 207 (1992).
- [17] G. Buchalla, A. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [18] M. Suzuki, Phys. Rev. D **33**, 676 (1986); C.-H. Chang and Y.-Q. Chen, Phys. Lett. B **284**, 127 (1992); Phys. Rev. D **46**, 3845 (1992); E. Braaten and T.C. Yuan, Phys. Rev. Lett. **71**, 1673; Phys. Rev. D **51**, 4819 (1995); E. Braaten, K. Cheung, and T. C. Yuan, Phys. Rev. D **48**, 4230 and R5049 (1993); R. Jaffe and L. Randall, Nucl. Phys. bf B421, 79 (1994).
- [19] T.C. Yuan, Phys. Rev. D **50**, 5664 (1994); K. Cheung and T.C. Yuan, Phys. Rev. D **53**, 1232 (1996); A.P. Martynenko and V.A. Saleev, Phys. Rev. D **53** 6666 (1996). They studied the $b \rightarrow B_c$ fragmentation, but one can read off the $c \rightarrow D_s$ fragmentation from their results.
- [20] R.L. Jaffe, Phys. Rev. D **15**, 267, 281 (1977).